

Rochester Brain and Spine

Dr Seth Zeidman

Emily Lapp PA-C Nathaniel Brochu PA-C Elizabeth Jefferson PA-C Cynthia Larson PA-C

400 Red Creek Drive, Suite 120; Rochester NY 14623

(585) 334-5560

Discharge Instructions

Cervical Laminectomy/Foramintomy/Discectomy

- **Medication:** You may use Tylenol or Ibuprofen as needed for pain. Narcotic pain medication may be dispensed and used on an as needed basis. Prescriptions for pain medications will be provided for up to 4 weeks after surgery. *After 4 weeks, you will need to have all pain medications managed by your primary care doctor or pain management doctor.*
- **Activity/Restriction:** You should walk daily, gradually increasing your time and distance as tolerated. Strict bed rest is not recommended as it will contribute to your stiffness and pain. Avoid repetitive reaching above your shoulder level. You may go up and down stairs as tolerated.

You should wear your cervical collar whenever you are out of bed until you are seen in the office for your first appointment [2 weeks after surgery]. You may remove your collar for showering, sleeping and eating. If you have any issues with your brace – contact Southern Cross Medical Supply at 585-334-5530.

Lifting is not to exceed 10 pounds. A gallon of milk is 8-9 pounds. You are to avoid strenuous activity.

You may begin to drive when you feel comfortable, about 3-4 weeks after surgery, provided that you have weaned yourself off of the narcotic pain medication and are out of the brace.

- **Wound Care:** You may remove the dressing over your incision(s) the day after you get home and leave the incision open to air. Keep the wound as dry as possible. You may also shower the day after you get home as well. Make sure to cover your incision(s) when in the shower [we recommend covering your incision(s) with Saran wrap and paper/silk tape] for the first week. You will have steri-strips [pieces of tape] over your incision(s). These will fall off on their own – about 10-14 days after surgery. Do NOT apply any lotions or creams to the incision [this includes Antibacterial cream, Neosporin or any scar creams]. No tub baths, pools or hot tubs until the incision(s) are well-healed [about 6-8 weeks].
- **General:** Make sure to drink plenty of fluids and use a stool softener after surgery. We recommend that you start taking Colace 100mg twice a day starting 5 days before your surgery and then increasing the Colace to 200mg twice a day after surgery. If you start to have diarrhea – contact the office and we will adjust the dose of the Colace.

Please call the office if you develop any of the following symptoms: fevers, chills, difficulty with urination or bowel movements [despite medications], persistent nausea or vomiting, severe pain not relieved with pain medications [some pain is to be expected], any redness or drainage from the incision. REMEMBER – smoking greatly reduces tissue healing. If you smoke, there is no better time to quit than now!